

# PHYTOCHEMICAL PROFILING OF PURNAJIWA FRUIT (Kopsia arborea Blume.) EXTRACTS AND ANTIBACTERIAL ACTIVITY AGAINST METHICILLIN-RESISTANT

Staphylococcus aureus (MRSA)

Maria Malida Vernandes Sasadara<sup>1\*</sup>, Erna Cahyaningsih<sup>1</sup>, Putu Era Sandhi Kusuma Yuda<sup>1</sup>, Ni Luh Putri Apriliani<sup>1</sup>, Ni Kadek Dwi Purnama Dewi<sup>1</sup>, Ni Luh Kade Arman Anita Dewi<sup>1</sup>

<sup>1</sup>Faculty of Pharmacy, Mahasaraswati University Denpasar \*Email Corresponding: mariasasadara@unmas.ac.id

Submitted: July 24, 2025 Revised: August 29, 2025 Accepted: September 2, 2025

### **ABSTRACT**

Purnajiwa (Kopsia arborea Blume.) contains diverse phytochemicals with notable antibacterial potential. This study aimed to characterize and quantify the phytochemicals of Kopsia arborea fruit extracts obtained by maceration and Soxhlet extraction with methanol and evaluate their antibacterial activity against MRSA through phytochemical screening, GC-MS analysis, spectrophotometric quantification of flavonoids, alkaloids, and phenolics, and disc diffusion assay at 100 µg/mL concentration. The findings indicated that the Soxhlet extraction produced a superior yield (19.47 ± 0.58%) compared to maceration (11.09 ± 0.65%). Phytochemical analysis revealed the presence of alkaloids, flavonoids, phenolics, tannins, saponins, and triterpenoids, with no qualitative differences between the two extracts. Quantitative analysis demonstrated higher concentrations of alkaloids, flavonoids, and phenolics in the Soxhlet extract, with values of  $88.58 \pm 3.76$ ,  $46.50 \pm 1.04$ , and  $57.87 \pm 0.44$  $\mu g/mL$ , respectively, compared to the maceration extract (50.46  $\pm$  3.86, 26.22  $\pm$  0.27, and  $30.49 \pm 0.31 \,\mu\text{g/mL}$ , respectively). GC-MS analysis identified 13 alkaloid compounds in the Soxhlet extract and 12 in the maceration extract. Antibacterial assays revealed that the mean inhibition zone diameter against MRSA was  $11.69 \pm 0.28$  mm for the Soxhlet extract and 12.61± 0.27 mm for the maceration extract, respectively. In conclusion, Soxhlet extraction yielded higher concentrations of alkaloids, phenolic compounds, and flavonoids; however, GC-MS analysis revealed that the macerated extract exhibited a higher AUC of alkaloid compounds than that of Soxhlet extraction. Moreover, the macerated extract demonstrated superior antibacterial activity, indicating that maceration has greater potential for development as an antibacterial agent than Soxhlet extraction.

Keywords: Kopsia arborea, maceration, Soxhlet extraction, phytochemical profiling, MRSA

## INTRODUCTION

Antibiotic resistance is a global problem that requires critical treatment. Cases of bacterial resistance are rapidly developing compared to the pace at which new antibiotics are developed. Antibiotic resistance in bacteria is a natural phenomenon and a form of bacterial adaptation to antimicrobial agents (Chandra *et al.*, 2017). The rapid development of bacterial resistance has become a concern in the medical world, prompting the search for alternative treatments or the development of new drug compounds to fight resistant bacteria. Research on alternative antimicrobials has been conducted intensively (Havenga *et al.*, 2019). Plants are a source of antimicrobial agents. Various studies have shown the effectiveness of plant metabolite compounds in fighting bacteria. Compounds commonly found in plants, such as quinine, alkaloids, lectins, polypeptides, flavonoids, manganese, terpenoids, essential oils, and tannins, have been shown to exhibit antibacterial activity (AlSheikh *et al.*, 2020; Chandra *et al.*, 2017).

Purnajiwa (*Kopsia arborea* Blume.) is a plant with potential antimicrobial properties. *Kopsia arborea* Blume is a species of the genus *Kopsia* (family *Apocynaceae*). *Kopsia* is known as one of the plants that produces alkaloid compounds. *Kopsia* species generally contain highly potent indole alkaloid compounds with a wide range of bioactivities (Chen *et al.*, 2020; Hop & Son, 2022; Wong *et al.*, 2021). Alkaloid compounds in *Kopsia* generally have a unique skeleton with significant bioactivity (Kogure *et al.*, 2012). Research on the ethanol extracts of *Kopsia arborea* Blume. shows the presence of several alkaloids, such as aspidospermidin, kopsinine, quebrachamin, and tabersonin (Ariati *et al.*, 2024).

Various studies have shown the effectiveness of alkaloid compounds as antibacterials. Alkaloid compounds naturally have inhibitory activity against Methicillin-resistant *Staphylococcus aureus* (MRSA). The indole alkaloid compounds dionemycin and 6-CH<sub>3</sub>O-7,7'-dichorochromopyrrolic acid, derived from *Streptomyces* sp., demonstrated inhibition against six strains of MRSA isolated from humans and pigs, with MIC in the range of 0.5–2μg/mL (Song *et al.*, 2020). Other alkaloid compounds include isoconcuressine and N-formylconessimine, isolated from the seeds of *Holarrhena antidysenteriaca* Wall. ex A. DC also shows inhibition against MRSA (Zhou *et al.*, 2017). Myoporumine A and myoporumine B, alkaloids isolated from *Myoporum bontioides* A. Gray, strongly inhibit MRSA with an MIC of 6.25μg/mL (Dong *et al.*, 2018).

Various studies have proven the antibacterial activity of various plants, which is produced by a variety of phytochemical compounds. This study aimed to identify and quantify the secondary metabolite components of Kopsia arborea Blume and evaluate its antibacterial activity against methicillin-resistant Staphylococcus aureus., which were extracted using two different methods: maceration and Soxhlet extraction. Both methods are conventional extraction methods commonly used to extract various secondary metabolite compounds, especially alkaloids. The antibacterial activity of a plant is attributed to its phytochemical compounds. Extraction techniques strongly influence the diversity and concentration of phytochemicals obtained from plant materials (Candra et al., 2021). In addition, variations in extraction techniques can also affect the interaction between solvents and soluble compounds, particularly those with similar polarity properties (Putri et al., 2022). Koçancı et al. (2022) proposed that Soxhlet extraction is a commonly used technique for extracting total alkaloids and identifying different types of alkaloid compounds. This method is also known to produce high extract yields with more economical solvent use and a relatively short duration (Marjoni 2016). Wirawan et al. (2023) showed that maceration can produce extracts with a wide range of phytochemical ingredients, such as flavonoids, phenols, alkaloids, and tannins. Maceration is a simple and effective method for maintaining the stability of heat-sensitive compounds, thereby minimizing the risk of damage to phytochemical compounds (Mukhriani, 2014; Rahman et al., 2017; Sekali et al., 2020). Several studies have also shown a correlation between antioxidant and antibacterial activities (Ispiryan et al., 2024; Todorovic et al., 2017).

This study presents a comparative evaluation of phytochemical content and antibacterial activity of *Kopsia arborea* fruit extracts obtained through maceration and Soxhlet methods, specifically targeting Methicillin-Resistant *Staphylococcus aureus* (MRSA). This study is among the first to directly contrast the influence of extraction techniques on both the phytochemical profile and antibacterial efficacy of purnajiwa (*Kopsia arborea* Blume.) against MRSA.

## RESEARCH METHODS

## **Sample Collection**

The harvesting of purnajiwa (*Kopsia arborea* Blume) fruit is collected in Denpasar, Bali (8°40'16.8"S 115°14'02.2"E). The fruit used was a ripe blackish-purple fruit. The sample was determined at the Mathematics and Natural Sciences Laboratory Unit, Faculty of Tarbiyah and Teacher Training, IAIN Syekh Nurjati Cirebon (27/In.08/LB.1.1/PP.009/12/2023). The samples were sorted, washed, dried, cut, and oven-dried at 50°C for 3 days until dry. The dry sample was then blended until a fine powder was obtained.

## **Extraction**

The sample powder was extracted by maceration and Soxhlet extraction using methanol with a sample and solvent ratio of 1:15. Maceration was carried out for 24 hours with periodic stirring. Soxhlet extraction was performed using a Soxhlet apparatus for 8 hours. Filtrates from both maceration methods were concentrated using a vacuum rotary evaporator at a controlled temperature of 40°C. Crude extracts obtained using both extraction methods were used in subsequent studies. The final result obtained was weighed and compared with the mass of the material used, calculated as a percentage of the extract yield.

% Yield =  $\frac{\text{final weight of extract}}{\text{Initial weight of extract}} \times 100\%$ 

# **Phytochemical Screening**

Phytochemical screening was performed as a preliminary identification of phytochemicals using standard colour reactions in test tubes. Screening was performed for compounds of the alkaloid group, flavonoids, phenols, tannins, quinones, saponins, steroids, and triterpenoids using a method described by Dubale *et al.* (2023).

The phytochemical content in crude extracts was also identified using the GC-MS Agilent Technologies 8860 GC system, 5977B GC/MSD (Agilent, USA) instrument, which is equipped with a column HP-5MS UI (30 m in length  $\times$  0.250 mm in diameter  $\times$  0.25  $\mu m$  in film). GC-MS spectroscopic detection was performed using an electron ionization system (70eV) with a range of 50–300 m/z. The temperature of the injector was 250 °C in the splitless mode. Helium was used as the carrier gas at a flow rate of 1 mL/min. The volume of the injected sample was 2  $\mu L$ . The initial temperature was set at 60–100°C with a rise rate of 4°C/min, and then increased to 290°C with a rise rate of 10°C/min. The presence of chemical compounds in the sample was characterized by the appearance of peaks in the chromatogram. The phytochemical compounds in the sample were identified by comparing the mass spectrum of the detected components with the spectral mass data of the components in the National Institute of Standards and Technology (NIST) library with a percentage similarity index of at least 60%. In this study, the NIST17 library was used. Data were analyzed using Agilent MassHunter Qualitative Analysis Navigator B.08.00 and GCMS 5977B software.

# **Quantification of Total Alkaloids**

The total alkaloid content was quantified using UV-VIS Spectrophotometry (Shimadzu, Japan) with caffeine as the standard. Linear regression was obtained from the calculation of the standard concentrations of caffeine (10, 15, 20, and 25 ppm) to the absorbance at a maximum wavelength of 270 nm. A total of 2 mL of extract solution with a concentration of 1000 ppm (in methanol) and a standard solution of caffeine were added with a pH phosphate buffer of 4.7 mL and a bromocresol green (BCG) solution of 2 mL, then extracted three times using 3 mL of chloroform. The chloroform phase was then evaporated using a water bath and dissolved in 10 mL of chloroform. The solution was then separated using a separate funnel, and the chloroform part was collected. The absorbance was measured at the maximum wavelength of caffeine (Karim et al., 2022). The determination was repeated thrice.

#### **Total Phenol Quantification**

Total phenol quantification was performed using UV-VIS Spectrophotometry (Shimadzu, Japan). A calibration curve was obtained using a gallic acid standard (0–60  $\mu$ g/mL). A total of 50  $\mu$ g/mL of extract and 1.6 mL of gallic acid were added to 0.2 mL of Folin-Ciocalteu reagent and mixed evenly for 3 minutes. Subsequently, 0.2 mL of 10% (b/b) sodium carbonate solution was added and left for 30 minutes at room temperature. Absorbance was measured at 760 nm for total phenolate content and 640 nm for total tannin compounds (Do *et al.*, 2014; McDonald *et al.*, 2001; Roghini & Vijayalakshmi, 2018). The determination was repeated three times.

#### **Total Quantification of Flavonoids**

Total flavonoid quantification was performed using UV-VIS Spectrophotometry (Shimadzu, Japan) with quercetin as the standard. Quercetin concentrations of 0– $100 \,\mu g/mL$  in methanol were used to generate a calibration curve for the total phenolate content. A total of 2.0 mL of extract was diluted in methanol to a concentration of  $100 \,\mu g/mL$  and quercetin was added to 0.1 mL solution of aluminum chloride 10% b/b and solution potassium acetate 0.1 mM. The mixture was left for 30 minutes at room temperature. Absorbance was measured at 415 nm (Do *et al.*, 2014). The determination was repeated thrice.

# **Antibacterial Activity Testing**

The antibacterial activity against MRSA was determined using the agar diffusion method with Mueller-Hinton Agar. Bacterial cultures were obtained from the Kerthi Bali Sadhajiwa Health Laboratory Center, Bali (UPTD). Balai Laboratorium Kesehatan Kerthi Bali Sadhajiwa Provinsi Bali). The bacteria were suspended at a bacterial concentration of  $1.5 \times 108$  CFU/mL, equivalent to the standard turbidity of Mc. Farland solution of 0.5, measured with Mc. Farland densitometer (DEN-1 Grant Instruments, England), and grown on a petri dish. The negative control used was dimethyl sufoxide (DMSO 5%), and the positive control was an antibiotic clindamycin 2  $\mu$ g (MRSA) disc. The test solution was prepared by macerating and extracting the fruit of *Kopsia arborea* Blume using Soxhlet extraction. suspended in 5% DMSO until a concentration of 100 mg/mL was obtained. The test was replicated three times and incubated at  $35-37^{\circ}$ C for 18 hours. The inhibition zone was observed, measured, and expressed in mm, excluding the disc size (6 mm).

## **Data Analysis**

The data obtained in this study were statistically analyzed using IBM SPSS Statistics (version 25) with a 95% confidence level. The total values of alkaloids, flavonoids, phenolics, and inhibitory zone diameters in both extracts were compared using an independent t-test to determine the differences in values between the Soxhlet and maceration extracts.

#### RESULTS AND DISCUSSION

Purnajiwa (*Kopsia arborea* Blume.) is a medicinal plant used in Indonesia. In this study, phytochemical and antibacterial activities were evaluated and compared for purnajiwa (*Kopsia arborea* Blume) fruit extracted using the Soxhlet and maceration methods. Qualitative phytochemical screening showed no difference in phytochemical content between the Soxhlet and maceration extracts, suggesting that both methods extracted the same type of metabolite (Table 1). Purnajiwa (*Kopsia arborea* Blume.) The fruit extracts contained alkaloids, flavonoids, phenolics, tannins, saponins, and triterpenoids. The results of this study are supported by previous research demonstrating that both maceration and Soxhlet extraction methods can isolate phytochemicals of similar types (Candra *et al.*, 2021; Putri *et al.*, 2024).

The higher extraction yield achieved by Soxhlet extraction compared to maceration in this study can be attributed to continuous hot solvent cycling, which enhances solvent penetration and compound diffusion from the plant matrix. The maceration method relies on passive diffusion and often requires an extended time to reach equilibrium, whereas Soxhlet extraction continuously replenishes fresh solvent over the sample, thereby increasing the mass transfer efficiency (Sun *et al.*, 2025). Several studies have reported similar findings, highlighting that the yields from Soxhlet extraction were markedly higher than those obtained via maceration (Bitwell *et al.*, 2023). These results also support the phytochemical quantification results of this study (Figure 1), which showed that Soxhlet extraction produces greater concentrations of alkaloids, phenolics, and flavonoids, owing to its more efficient extraction dynamics under heat and solvent circulation.

**Table I.** Yield and Phytochemical Screening Results on Maceration and Soxhlet Extraction Extracts of Purnajiwa Fruit (*Kopsia arborea* Blume.)

Donomotons	Result		
Parameters	Soxhlet Extraction	Maceration	
Extract yields	$19.47 \pm 0.58 \%$	$11.09 \pm 0.65 \%$	
Phytochemical screening			
Alkaloid (Dragendorf)	+	+	
Alkaloids (Mayer)	+	+	
Flavonoids	+	+	
Phenolic	+	+	
Tannins	+	+	
Quinone	-	-	
Saponins	+	+	
Steroids	-	-	
Triterpenoids	+	+	

Caption: (+) detected; (-) Undetectable




Figure 1. Concentrations of alkaloids, flavonoids, and phenolics in the maceration and Soxhlet extracts of the Purnajiwa fruit (*Kopsia arborea*). The asterisk (\*) indicates a significant difference (p < 0.05) between the Soxhlet and maceration extract.

The quantification of alkaloids, flavonoids, and phenolic compounds in both extracts showed differences in compound levels (Figure 1). Extraction by the Soxhlet method yielded higher alkaloids, flavonoids, and phenolic compounds than maceration. Other studies, such as those on *Urtica dioica*, have shown that Soxhlet extraction is capable of producing a higher total amount of phenols and flavonoids than maceration (Horozić *et al.*, 2023). The application of the Soxhlet technique for extraction can maintain higher temperatures and continuous solvent circulation, which improves the extraction efficiency of polar and heat-sensitive compounds. Maaiden *et al.* showed that Soxhlet extraction resulted in a higher polyphenol extract than maceration (Maaiden *et al.*, 2022).

The difference in extraction efficiency between the maceration and Soxhlet extraction methods is estimated to be influenced by several factors, such as solvent dynamics, temperature control, and the continuous extraction process. In Soxhlet extraction, a solvent circulation mechanism allows the thorough dissolution of bioactive compounds from the plant matrix until it is completely saturated, thus facilitating a more efficient extraction process (Horozić *et al.*, 2023). This is in contrast to maceration, which relies on soaking plant material

in solvents at room temperature, often leading to incomplete extraction (Rachmaniah *et al.*, 2021). In addition, heating during Soxhlet extraction can increase the solubility and diffusion of phenolic compounds into solvents (Iftikhar *et al.*, 2022). However, it should be noted that heating can also degrade phytochemical compounds. In addition, the special design of the Soxhlet apparatus allows repeated contact between plant material and fresh solvents, thereby improving the mass transfer and dissolution of bioactive molecules from the plant matrix (Ghanimi *et al.*, 2022; Raghu & Velayudhannair, 2023).

Plants of the genus *Kopsia* are known to produce alkaloid compounds. In this study, alkaloids in both extracts were specifically identified using GC-MS. Table II lists the alkaloid compounds identified in both extracts. There were 13 compounds in the Soxhlet extraction and 12 compounds in the maceration extract. However, the percent area under the curve (AUC) value of these compounds showed that maceration extracts (14.85%) had a larger area value of alkaloid compounds than Soxhlet extraction (7.05%).

In this study, Soxhlet extraction yielded a higher total alkaloid content in the quantitative assays, whereas GC-MS analysis indicated a larger percentage area of alkaloids in the macerated extract. This discrepancy can be explained by the inherent limitations of GC-MS, which primarily detects volatile and thermally stable compounds. Certain alkaloids present in the Soxhlet extract may not have been detected because they are less volatile or decompose upon heating, even though their total concentrations are higher. Moreover, the area under the curve (AUC) values generated by GC-MS were calculated relative to the total number of peaks detected.

Aspidospermidine is an alkaloid compound with the largest area value (5.80%) that appears in the maceration extraction. Aspidospermin is an alkaloid compound widely found in the plant genus *Aspidosperma*. This compound has a distinctive structural framework. Some studies have shown the various pharmacological activities of aspidospermidin, such as aspidospermidin-17-ol from *Monascus purpureus*, with bioactive activities, including antiviral, anti-inflammatory, immunomodulatory, and anticancer activities (Teiba *et al.*, 2024).

**Table II.** Alkaloid Compounds Identified on Chromatograms (GC-MS) in Macerated and Soxhlet Extraction Extracts of the Ripe Fruit (*Kopsia arborea Blume.*)

		Extraction Method			
No.	Compound Name	Soxhletasi		Maceration	
		RT	AUC	RT	AUC
1.	Pyridine, 3-ethyl-	4.223	0.92	3.956	0.64
2.	2H-Pyrazole-3-carboxylic acid, 2-methyl-	-	-	7.838	0.09
3.	Ethanone, 1-(1-ethyl-1H-pyrazol-yl)-	12.765	0.76	-	-
4.	1H-Indole-5-ol	-	-	13.762	0.09
5.	[1,2,5]Oxadiazolo[3,4-b][1,4]diazocin- 5,7(4H,6H)-dione, 8,9 dihydro	14.223	0.11	-	-
6.	Aspidospermidine, 1,2-didehydro-,(5.alpha.,12.beta.,19.alpha.)-	-	-	21.726	1.87
7.	3,5-Dimethyl-1-[4-(1H-pyrrol-1-yl)phenyl]-1H-pyrazole	23.019	0.12	-	-
8.	Quebrachamine	-	_	23.939	1.71
9.	Aspidospermidine-3-carboxylic acid, 2,3-didehydro-, methyl ester,	-	-	25.003	5.80
10.	(5.alpha.,12.beta.,19.alpha.)- Aspidofractinine-3-carboxylic acid , methyl ester, (2.alpha.,3.alpha. ,5.alpha.)-	25.030	1.08	-	-
11.	4,6 -Dimethyl -2-pyrimidone	25.290	0.53	-	-

		Extraction Method			
No.	<b>Compound Name</b>	Soxhletasi		Maceration	
	- -	RT	AUC	RT	AUC
12.	Indeno[1,2-b]quinoline, 11-(4-	25.487	1.90	-	-
	pyridylmethylene)-				
13.	11-Benzyl-2-methyl-2,3,4,5,6,11-	25.792	0.26	-	-
	hexahydro-azocino[3,4-b]indole-1-one				
14.	1H-Indole, 3-(5-methyl-2-(pyridin-3-	26.032	0.11	-	-
	yl)thiazol-4-yl-				
15.	1,2-Benzisothiazole,3-(hexahydro-1H-	26.589	0.35	-	-
	azepin-1-yl)-, 1,1-dioxide				
16.	Aspidofractinine-3-carboxylic acid, methyl	26.828	0.39	25.071	1.49
	ester, (2.alpha.,3.alpha.,5.alpha.)-				
17.	11-Butyl-8-methoxy-11H-indolo[3,2-	-	-	25.209	1.42
	c]quinoline				
18.	Aspidofractinine-3-carboxylic acid 1-	-	-	26.843	0.73
	formyl-17-methoxy-, methyl ester,				
	(2.alpha.,3.beta.,5.alpha.)-				
19.	1-[3-Amino-4-(pyridin-4-yl)-5H,6H1-[3-	-	-	27.501	0.41
	Amino-4-(pyridin-4-yl)-5H,6Hthanone				
20.	Benzenamine, 4,4'-[2-(4-methoxyphenyl)-	-	-	27.690	0.60
	1H-imidazole-4,5-diyl]bis[N,N-dimethyl-				
21.	2-Phenyl-3-pyridin-4-yl-7-(pyridin-4	28.219	0.52	-	-
	ylmethylidene)-3,3a,4,5,6,7-hexahydro-				
	2H-indazole				
22.	Oxayohimban-16-carboxylic acid, 16,17-	28.966	0.22	29.168	0.97
	didehydro-19-methyl-, methyl e				
	ster,(19.alpha.,20.alpha.)-				




Figure 2. Chromatogram (GC-MS) on a soxhletation extract of the deceased fruit (Kopsia arborea Blume.)




Figure 3. Chromatogram (GC-MS) on the macerated extract of the deceased fruit (Kopsia arborea Blume.)




Figure 4. Results of antibacterial assay soxhletation (S) and maceration (M) extract of purnajiwa (Kopsia arborea Blume) fruit

The results of the measurement of the diameter of the inhibition zone from the extract of the purnajiwa fruit (*Kopsia arborea* Blume.) against the growth of MRSA (Methicillin-Resistant *Staphylococcus aureus*) bacteria are presented in Figure 4 and Table III. The antibacterial activity test showed that the macerated extract produced a larger inhibition zone than Soxhlet extraction. The results of this study show that the high total content of alkaloids, phenolics, and flavonoids in Soxhlet extract is not in line with the increase in antibacterial activity produced. It is estimated that certain compounds with antibacterial activity are degraded when the Soxhlet method is used, which changes their chemical structure and decreases their sensitivity to bacteria. In contrast, maceration at room temperature results in extracts with a lower total phytochemical content but a stable and intact composition of active compounds. Minor compounds or secondary metabolites with MRSA-inhibiting activity are likely to be better preserved during maceration, resulting in higher antibacterial effectiveness, even though the total number of compounds extracted is lower. In addition, maceration can maintain a natural balance of phytochemicals that favors synergistic effects.

Table III. Measurement Results of Inhibition Zone Diameter

Donlingtion	Inhibitory Zone Diameter (mm)			
Replication -	Soxhlet	Maceration	C-	
1	11.43	12.74	0	
2	11.98	12.28	0	
3	11.66	12.26	0	
Average ± SD	11.69±0.28	12.61±0.27*	0	

Remarks: SD = standard deviation; C- = Negative Control (5% DMSO Solution)

The actorists (\*) indicates a significant difference (p. 6.05) between the S.

The asterisk (\*) indicates a significant difference (p < 0.05) between the Soxhlet and maceration extract.

In this study, the inhibitory activity of purnajiwa (*Kopsia arborea B*lume.) extracts was classified as strong, with inhibition zone diameters ranging from 10–20 mm. These results indicate that the purnajiwa fruit extract demonstrates promising potential as a natural antibacterial agent, warranting further investigation to support its application in addressing antibiotic resistance. A recent comprehensive review highlighted that both flavonoids and alkaloids are among the most effective phytochemicals against MRSA, acting through mechanisms such as membrane disruption and efflux pump inhibition (Liang *et al.*, 2022).

Alkaloids, phenolics, and flavonoids contribute to antibacterial activity against MRSA through distinct but complementary mechanisms. Alkaloids disrupt bacterial cell membrane integrity and interfere with essential metabolic pathways, such as peptidoglycan synthesis, thereby weakening their structural stability (Prakoso *et al.*, 2020). Phenolic compounds exert broad-spectrum effects by forming complexes with bacterial components, altering membrane permeability, and disrupting the biosynthesis of structural components (Elmaidomy *et al.*, 2023; Rahayu *et al.*, 2020). Flavonoids further enhance anti-MRSA efficacy by modulating key target proteins, such as PBP2a, increasing susceptibility to β-lactam antibiotics, reducing membrane fluidity, and impairing bacterial energy production (Jieputra *et al.*, 2024; Rahayu *et al.*, 2020).

#### **CONCLUSION**

This study demonstrates that the extraction method significantly affects the yield, phytochemical content, and antibacterial activity of the Purnajiwa (*Kopsia arborea* Blume) fruit extract. The Soxhlet method resulted in a higher yield and total content of alkaloids, phenolics, and flavonoids than maceration, supported by the identification of 13 alkaloid compounds through GC-MS analysis. However, macerated extracts consistently showed larger diameters of inhibition zones against Methicillin-Resistant *Staphylococcus aureus* (MRSA), indicating better antibacterial activity.

# ACKNOWLEDGEMENT

Thanks to the Faculty of Pharmacy, Mahasaraswati University, Denpasar, for funding this research (1119.13/E.003/FF-UNMAS/XII/2024)

#### REFERENCES

AlSheikh, H. M. Al., Sultan, I., Kumar, V., Rather, I. A., Al-Sheikh, H., Tasleem Jan, A., & Haq, Q. M. R. (2020). Plant-Based Phytochemicals as Possible Alternatives to Antibiotics in Combating Bacterial Drug Resistance. *Antibiotics (Basel, Switzerland)*, 9(8). https://doi.org/10.3390/antibiotics9080480

Ariati, P. E. P., Wirawan, I. G. P., Sasadara, M. M. V, Jawi, I. M., Sunyamurthi, I. G. N. A., & Wijaya, I. N. (2024). Phytochemical Profiling of Balinese Alkaloid-Source Plant Purnajiwa (*Kopsia arborea* Blume. and *Euchresta horsfieldii* (Lesch.) Benn.). *Tropical Journal of Natural Product Research* (*TJNPR*), 8(5), 7089–7095. https://doi.org/10.26538/tjnpr/v8i5.5

- Bitwell, C., Indra, S. Sen, Luke, C., & Kakoma, M. K. (2023). A Review of Modern and Conventional Extraction Techniques and Their Applications for Extracting Phytochemicals from Plants. *Scientific African*, 19, e01585. https://doi.org/10.1016/j.sciaf.2023.e01585
- Candra, L. M. M., Andayani, Y., & Wirasisya, D. G. (2021). Pengaruh Metode Ekstraksi Terhadap Kandungan Fenolik Total dan Flavonoid Total Pada Ekstrak Etanol Buncis (*Phaseolus Vulgaris* L.). *Pilar MIPA*, 16(3), 397–405. https://doi.org/10.29303/jpm.v16i3.2308
- Chandra, H., Bishnoi, P., Yadav, A., Patni, B., Mishra, A. P., & Nautiyal, A. R. (2017). Antimicrobial Resistance and Alternative Resources with Special Emphasis on Plant-Based Antimicrobials: A Review. *Plants (Basel, Switzerland)*, 6(2). https://doi.org/10.3390/plants6020016
- Chen, X. D., Hu, J., Li, J. X., & Chi, F. S. (2020). Cytotoxic Monoterpenoid Indole Alkaloids from the Aerial Part of *Kopsia arborea*. *Journal of Asian Natural Products Research*, 22(11), 1024–1030. https://doi.org/10.1080/10286020.2019.1680646
- Do, Q. D., Angkawijaya, A. E., Tran-Nguyen, P. L., Huynh, L. H., Soetaredjo, F. E., Ismadji, S., & Ju, Y. H. (2014). Effect of Extraction Solvent on Total Phenol Content, Total Flavonoid Content, and Antioxidant Activity of *Limnophila aromatica*. *Journal of Food and Drug Analysis*, 22(3), 296–302. https://doi.org/10.1016/j.jfda.2013.11.001
- Dong, L.-M., Huang, L.-L., Dai, H., Xu, Q.-L., Ouyang, J.-K., Jia, X.-C., Gu, W.-X., & Tan, J.-W. (2018). Anti-MRSA Sesquiterpenes from the Semi-Mangrove Plant *Myoporum bontioides* A. Gray. *Marine Drugs*, *16*(11). https://doi.org/10.3390/md16110438
- Dubale, S., Kebebe, D., Zeynudin, A., Abdissa, N., & Suleman, S. (2023). Phytochemical Screening and Antimicrobial Activity Evaluation of Selected Medicinal Plants in Ethiopia. *Journal of Experimental Pharmacology*, 15, 51–62. https://doi.org/10.2147/JEP.S379805
- Elmaidomy, A. H., Mohamad, S. A., Abdelnaser, M., Yahia, R., Mokhtar, F. A., Alsenani, F., Badr, M. Y., Almaghrabi, S., Altemani, F. H., Alzubaidi, M. A., Saber, E. A., Elrehany, M. A., Abdelmohsen, U. R., & Sayed, A. M. (2023). *Vitis vinifera* Leaf Extract Liposomal Carbopol Gel Preparation's Potential Wound Healing and Antibacterial Benefits: In Vivo, Phytochemical, and Computational Investigation. *Food & Function*, 14(15), 7156–7175. https://doi.org/10.1039/d2fo03212k
- Ghanimi, R., Ouhammou, A., Atki, Y. E., Bouchari, M. E. H., & Cherkaoui, M. (2022). The Antioxidant Activities of Ethanolic, Methanolic, Ethyl Acetate, and Aqueous Extracts of the Endemic Species, *Lavandula mairei* Humbert (A Comparative Study Between Cold and Hot Extraction). *Ethiopian Journal of Health Sciences*, 32(6). https://doi.org/10.4314/ejhs.v32i6.21
- Havenga, B., Ndlovu, T., Clements, T., Reyneke, B., Waso, M., & Khan, W. (2019). Exploring the Antimicrobial Resistance Profiles of WHO Critical Priority List Bacterial Strains. *BMC Microbiology*, *19*(1), 1–16. https://doi.org/10.1186/s12866-019-1687-0
- Hop, N. Q., & Son, N. T. (2022). A Comprehensive Review on Phytochemistry and Pharmacology of Genus Kopsia: Monoterpene Alkaloids - Major Secondary Metabolites. RSC Advances, 12(30), 19171–19208. https://doi.org/10.1039/d2ra01791a
- Horozić, E., Begić, S., Zukić, A., Alibašić, H., & Briga, M. (2023). Effect of Extraction Technique on the Content of Bioactive Components and Antioxidant Activity of Aqueous Extracts of Fresh and Dried Nettle (*Urtica dioica L.*). *Technol. Acta*, *15*(2). https://doi.org/10.51558/2232-7568.2022.15
- Iftikhar, S., Hussain, K., Saleem, M., Shehzadi, N., Saghir, F., Shaukat, A., Naheed, S., & Siddique, S. (2022). Impact of Extraction Mode on Phytoconstituents and Antidiabetic Activity of Extracts of *Calotropis procera* (Aiton) Dryand. *Journal of Pharmacy & Pharmacognosy Research*, 10(1), 30–38.
- Ispiryan, A., Atkociuniene, V., Makstutiene, N., Sarkinas, A., Salaseviciene, A., Urbonaviciene, D., Viskelis, J., Pakeltiene, R., & Raudone, L. (2024). Correlation

- between Antimicrobial Activity Values and Total Phenolic Content/Antioxidant Activity in *Rubus idaeus* L. *Plants*, *13*(4). https://doi.org/10.3390/plants13040504
- Jieputra, A., Purwanta, M., Mustika, A., & Retnowati, W. (2024). In Vitro Antimicrobial Activity Test of *Zingiber officinale* Var. Rubrum Rhizome Extract Against Methicillin-Sensitive *Staphylococcus aureus* (MSSA) and Methicillin-Resistant *Staphylococcus aureus* (MRSA). *Juxta Jurnal Ilmiah Mahasiswa Kedokteran Universitas Airlangga*, 15(1), 57–63. https://doi.org/10.20473/juxta.v15i12024.57-63
- Karim, Abd., Adnan, J., & Irmawati. (2022). Penentuan Kadar Alkaloid Total Ekstrak Etanol Daun Ungu (*Graptophyllum pictum* L.) dengan Metode Spektrofotometri Uv-Vis. *Farmasi Pelamonia*, 42–47.
- Koçancı, F. G., Dolanbay, S. N., & Aslım, B. (2022). Comparison of Three Different Protocols of Alkaloid Extraction from *Glaucium corniculatum* Plant. *International Journal of Secondary Metabolite*, 9(1), 43-51. http://dx.doi.org/10.21448/ijsm.980171
- Kogure, N., Suzuki, Y., Wu, Y., Kitajima, M., Zhang, R., & Takayama, H. (2012). Chemical Conversion of Strychnine into Kopsiyunnanine-I, A New Hexacyclic Indole Alkaloid from Yunnan *Kopsia arborea*. *Tetrahedron Letters*, *53*(48), 6523–6526. https://doi.org/10.1016/j.tetlet.2012.09.078
- Liang, M., Ge, X., Xua, H., Ma, K., Zhang, W., Zan, Y., Efferth, T., Xue, Z., & Hua, X. (2022). Phytochemicals with Activity Against Methicillin-Resistant *Staphylococcus aureus*. *Phytomedicine*, 100, 154073.
- Maaiden, E. E., Bouzroud, S., Nasser, B., Moustaid, K., Mouttaqi, A. E., Ibourki, M., Boukcim, H., Hirich, A., Kouisni, L., & Kharrassi, Y. E. (2022). A Comparative Study between Conventional and Advanced Extraction Techniques: Pharmaceutical and Cosmetic Properties of Plant Extracts. *Molecules*, 27(7), 2074. https://doi.org/10.3390/molecules27072074
- Marjoni, Mhd. R. (2016). *Dasar-Dasar Fitokimia untuk Diploma III Farmasi* (T. Ismail, Ed.). CV. Trans Info Media.
- McDonald, S., Prenzler, P. D., Antolovich, M., & Robards, K. (2001). Phenolic Content and Antioxidant Activity of Olive Extracts. *Food Chemistry*, 73(1), 73–84. https://doi.org/10.1016/S0308-8146(00)00288-0
- Mukhriani. (2014). Ekstraksi, Pemisahan Senyawa, dan Identifikasi Senyawa Aktif. *Jurnal Kesehatan*, VII(2/2014).
- Prakoso, Y. A., Rini, C. S., Rahayu, A., Sigit, M., & Widhowati, D. (2020). Celery (*Apium graveolens*) as a Potential Antibacterial Agent and Its Effect on Cytokeratin-17 and Other Healing Promoters in Skin Wounds Infected With Methicillin-Resistant *Staphylococcus aureus*. *Veterinary World*, *13*(5), 865–871. https://doi.org/10.14202/vetworld.2020.865-871
- Putri, C. N., Rahardhian, M. R. R., & Ramonah, D. (2022). Pengaruh Metode Ekstraksi Terhadap Kadar Total Fenol dan Total Flavonoid Ekstrak Etanol Daun Insulin (Smallanthus sonchifolius) serta Aktivitas Antibakteri Terhadap Staphylococcus aureus. JPSCR: Journal of Pharmaceutical Science and Clinical Research, 7(1), 15. https://doi.org/10.20961/jpscr.v7i1.43465
- Putri, V. M., Azzahra, F., Utami, A., & Astuti, F. (2024). Perbandingan Rendemen dan Kandungan Kimia Secara Kualitatif Ekstrak Daun Alpukat (*Persea americana*) Berdasarkan Perbedaan Metode Ekstraksi. *Jurnal Kesehatan Bakti Tunas Husada: Jurnal Ilmu Keperawatan, Analis Kesehatan, dan Farmasi*.
- Rachmaniah, O., Spronsen, J. van, Verpoorte, R., & Witkamp, G.-J. (2021). The Effect of Acids on Alkaloid Yield in Pressurized Water Extraction of *Narcissus pseudonarcissus*. *Journal of Engineering and Technological Sciences*, 53(6), 210608. https://doi.org/10.5614/j.eng.technol.sci.2021.53.6.8
- Raghu, A., & Velayudhannair, K. (2023). Phytochemical Analysis and Antibacterial Potential of *Stevia Rebaudiana* (Bertoni, 1899) Leaf Extracts Against *Aeromonas* Species: Influence of Extraction Methods and Solvents in Aquaculture Applications. *Journal of*

- *Pure and Applied Microbiology*, *17*(4), 2352–2366. https://doi.org/10.22207/jpam.17.4.31
- Rahayu, A., Rini, C. S., Prakoso, Y. A., Palgunadi, B. U., & Munandar, M. A. (2020). Effects of *Sauropus androgynus* Extract and Its Combination with Ampicillin Against Methicillin-Resistant *Staphylococcus aureus*: An In Vitro Study. *International Journal of One Health*, 6(2), 128–133. https://doi.org/10.14202/ijoh.2020.128-133
- Rahman, A., Taufiqurrahman, I., & Edyson. (2017). Perbedaan Total Flavonoid antara Metode Maserasi dengan Sokletasi Pada Ekstrak Daun Ramania (*Bouea macrophylla* Griff). *Dentino*, *I*(1), 22–27.
- Roghini, R., & Vijayalakshmi, K. (2018). Phytochemical Screening, Quantitative Analysis of Flavonoids and Minerals in Ethanolic Extract of *Citrus paradisi*. *International Journal of Pharmaceutical Sciences and Research*, 9(11), 4859–4864. https://doi.org/10.13040/IJPSR.0975-8232.9(11).4859-64
- Sekali, E. E. K., Wartini, N. M., & Suhendra, L. (2020). Karakteristik Ekstrak Aseton Pewarna Alami Daun Singkong (*Manihot esculenta* C.) pada Perlakuan Ukuran Partikel Bahan dan Lama Maserasi. *Jurnal Ilmiah Teknologi Pertanian Agrotechno*, *5*(2):49. https://doi.org/10.24843/JITPA.2020.v05.i02.p02
- Song, Y., Yang, J., Yu, J., Li, J., Yuan, J., Wong, N.-K., & Ju, J. (2020). Chlorinated Bis-Indole Alkaloids from Deep-Sea Derived *Streptomyces* sp. SCSIO 11791 with Antibacterial and Cytotoxic Activities. *The Journal of Antibiotics*, 73(8), 542–547. https://doi.org/10.1038/s41429-020-0307-4
- Sun, S., Yu, Y., Jo, Y., Han, J. H., Xue, Y., Cho, M., Bae, S.-J., Ryu, D., Park, W., Ha, K.-T., & Zhuang, S. (2025). Impact of Extraction Techniques on Phytochemical Composition and Bioactivity of Natural Product Mixtures. *Frontiers in Pharmacology*, 16. https://doi.org/10.3389/fphar.2025.1615338
- Teiba, I. I., Mazrou, Y. S. A., Makhlouf, A. H., Nehela, Y., Mohamed, A. E., Abbas, A. M., Mamdouh, I., & El-Bilawy, E. H. (2024). Antibacterial Potential of Honeybee Venom and *Monascus purpureus* Extracellular Metabolites Against Multidrug-Resistant Pathogenic Bacteria. *Biology*, 14(1), 21. https://doi.org/10.3390/biology14010021
- Todorovic, V., Milenkovic, M., Vidovic, B., Todorovic, Z., & Sobajic, S. (2017). Correlation between Antimicrobial, Antioxidant Activity, and Polyphenols of Alkalized/Nonalkalized Cocoa Powders. *Journal of Food Science*, 82(4), 1020–1027. https://doi.org/10.1111/1750-3841.13672
- Wirawan, I. G. P., Sasadara, M. M. V., Jawi, I. M., Darmawati, I. A. P., Wijaya, I. N., Krisnandika, A. A. K., Sunyamurthi, I. G. N. A., & Wirya, I. G. N. A. S. (2023). Balinese Purnajiwa (Kopsia arborea Blume.) Extract Stimulates Male Rats' Sexual Behavior and Plasma Testosterone Level. Bali Medical Journal, 12(1), 1026–1032. https://doi.org/10.15562/bmj.v12i1.4163
- Wong, S. K., Yeap, J. S. Y., Tan, C. H., Sim, K. S., Lim, S. H., Low, Y. Y., & Kam, T. S. (2021). Arbolodinines A–C, Biologically-Active Aspidofractinine-Aspidofractinine, Aspidofractinine-Strychnan, and Kopsine-Strychnan Bisindole Alkaloids from *Kopsia arborea*. Tetrahedron, 78, 131802. https://doi.org/10.1016/j.tet.2020.131802
- Zhou, L.-N., Ge, X.-L., Dong, T.-T., Gao, H.-Y., & Sun, B.-H. (2017). Antibacterial Steroidal Alkaloids from *Holarrhena antidysenteriaca*. *Chinese Journal of Natural Medicines*, 15(7), 540–545. https://doi.org/https://doi.org/10.1016/S1875-5364(17)30080-8